

DEWORMING GUIDELINES

<u>MONTH</u>	<u>DRUG NAME</u>	<u>COMMON NAMES</u>	<u>REASONS</u>	<u>OTHER "TO DO's"</u>
JANUARY				
FEBRUARY	FENBENDAZOLE for 5 consecutive days	PANACUR POWERPAC	Panacur for 5 consecutive days at the larvacidal (double) dose is effective at killing all encysted larvae.	OR you can use STRONGID
MARCH				FECAL EXAM
APRIL	IVERMECTIN & PRAZIQUANTEL	EQUIMAX or ZIMECTERIN GOLD	This combination drug has recently been approved and is effective against all parasites, including TAPEWORMS.	
MAY				
JUNE	MOXIDECTIN	QUEST	DO NO USE in animals less than 6 months old.	
JULY				
AUGUST	PYRANTEL	STRONGID		
SEPTEMBER				FECAL EXAM
OCTOBER	IVERMECTIN & PRAZIQUANTEL	EQUIMAX or ZIMECTERIN GOLD	OR use straight IVERMECTIN (I.e. EQUELL)	
NOVEMBER				
DECEMBER	MOXIDECTIN	QUEST		

DEWORMING GUIDELINES

NOTES	Pyrantel	DAILY STRONGID 2X	IF using the DAILY STRONGID, then paste worm TWICE a year with EQUIMAX or EQUELL , spring and fall.	
-------	----------	--------------------------	---	--